

# Entropy of BTZ black hole and its spectrum by quantum radial geodesics behind horizons

V.V.Kiselev\*

*Russian State Research Center "Institute for High Energy Physics",  
Pobeda 1, Protvino, Moscow Region, 142281, Russia  
Fax: +7-0967-744937*

In the framework of thermal quantization of radial geodesics completely confined behind the horizons we calculate the entropy of BTZ black hole in agreement with the Bekenstein–Hawking relation. Particles in the BTZ black hole occupy the only quantum ground level. The quantization allow us to find a linear dependence of black hole mass versus its orbital momentum.

PACS numbers: 04.70.Dy

## I. INTRODUCTION

Recently, we have formulated a new framework for calculating the entropy of black hole by a quasi-classical description of particles moving at radial geodesics completely confined behind horizons of black hole [1, 2, 3, 4]. The thermodynamical assignment of such the motion is twofold. First, the trajectories are periodic in purely imaginary time, which is a direct indication of thermal equilibrium in terms of path integral for the partition function. The periodicity leads to a thermal quasi-classical quantization, so that the partition function is approximated by a leading term of sum over the discrete geodesics. Second, the macro-state of black hole is fixed by its mass and other several global charges as an electric charge or an orbital momentum, while the micro-state is composed of full set of quantum levels for each particle at confined geodesics. Therefore, the macro-state is not a pure quantum state, it is characterized by a density matrix obtained by summing up over the micro-states. Such the approach has been tested on standard examples of black holes, i.e. the Schwarzschild black hole, and Reissner–Nordstöm, Kerr–Newman ones [2]. The Hawking radiation [5] of Schwarzschild black hole by transitions between the levels of particle confined behind the horizon has been investigated in [4].

In present paper we expand the analysis to the Bañados–Teitelboim–Zanelli black hole [6].

The motion of massive particles in the gravitational field of BTZ black hole can be studied by the action given by the sum of gravity and particles terms

$$S = \frac{1}{16\pi G} \int \sqrt{-g} d^3x \left\{ R + \frac{2}{\ell^2} \right\} - \sum mc \int ds, \quad (1)$$

where  $R$  is the Ricci scalar of curvature,  $G$  is the gravitational constant with the dimension of length or inverse energy, the length  $\ell$  represents the cosmological constant of anti-de Sitter (AdS) 2+1 space-time, while the interval

of BTZ black hole is given by

$$d^2s = N^2 dt^2 - \frac{1}{N^2} d\rho^2 - \rho^2 \left( -\frac{4JG}{\rho^2} dt + d\phi \right)^2. \quad (2)$$

In (2)

$$N^2 = \frac{1}{\ell^2 \rho^2} (\rho^4 - 8GM\ell^2 \rho^2 + 16J^2 G^2 \ell^2), \quad (3)$$

whereas  $J$  is the orbital momentum of black hole.

The BTZ black hole has two horizons (nullifying the function  $N^2$ )

$$\rho_{\pm}^2 = 4GM\ell^2 \left( 1 \pm \sqrt{1 - \frac{J^2}{M^2 \ell^2}} \right), \quad (4)$$

determining two 1D ‘areas’

$$\mathcal{A}_{\pm} = 2\pi \rho_{\pm}. \quad (5)$$

The solution is the black hole, if  $\rho_{\pm}$  are real numbers, i.e.

$$|J| \leq M\ell. \quad (6)$$

In section II we study the BTZ black hole with  $J = 0$ . We in detail describe the introduction of appropriate coordinates for geodesics completely confined behind the horizon and the thermodynamical quantization, which results in the only ground level of particle in the field of BTZ black hole. Fixing the total energy of particles by the black hole mass, we derive the entropy satisfying the Bekenstein–Hawking law. The case of  $J \neq 0$  with two horizons is investigated in section III. Consistence of two maps behind the horizons leads to quantizing the ratio of horizon radii. The ratio has to run a finite set of values, since we introduce a winding number equal to a number of cycles per period in imaginary time, and it should be integer. The entropy of BTZ black hole in general case is reproduced. Section IV represents the quasi-classical spectrum of BTZ black hole: the mass is linear in  $J$ , and the area squared is also linear in  $J$ . Our conclusions are listed in section V.

---

\*Electronic address: kiselev@th1.ihep.su

## II. NO ROTATION: $J = 0$

At  $J = 0$  the metric reduces to

$$ds^2 = N_0^2 dt^2 - \frac{1}{N_0^2} dr^2 - r^2 d\phi^2, \quad (7)$$

with

$$N_0^2 = \frac{r^2 - r_+^2}{\ell^2}, \quad r_+^2 = 8GM\ell^2. \quad (8)$$

At radial geodesics

$$(\dot{r})^2 = N_0^4(1 - AN_0^2), \quad (9)$$

where the integral of motion  $A$  is determined by the total energy of particle,  $E$ , with the mass  $m$ ,

$$A = \frac{m^2}{E^2}. \quad (10)$$

The interval at the trajectory is expressed by

$$ds^2 = \frac{A}{1 - AN_0^2} dr^2 = \frac{\ell^2}{r_c^2 - r^2} dr^2, \quad (11)$$

where

$$r_c^2 = r_+^2 + \frac{\ell^2}{A}. \quad (12)$$

For the radial geodesics completely confined behind the horizon, we get

$$r_c^2 < r_+^2 \quad \Rightarrow \quad A < 0, \quad (13)$$

so that

$$ds^2 > 0, \quad \text{if } r: 0 < r < r_c < r_+. \quad (14)$$

At such causal geodesics

$$ds^2 = AN_0^4 dt^2 > 0, \quad \text{at } A < 0, \quad \Rightarrow \quad dt^2 < 0, \quad (15)$$

i.e. the time should run purely imaginary values. Such evolution in the imaginary time can be assigned to the thermodynamical character of motion. Namely, a period in imaginary time is equal to inverse temperature of thermodynamical ensemble in a thermal equilibrium. In order to fix the period let us to consider the Kruskal coordinates.

First, the ‘tortoise’ coordinate is determined by

$$r_* = \int \frac{dr}{N_0^2} = \frac{\ell^2}{2r_+} \ln \frac{r - r_+}{r + r_+}, \quad (16)$$

while the Kruskal coordinates are

$$\begin{cases} u = t - r_*, \\ v = t + r_*, \end{cases} \quad \Rightarrow \quad \begin{cases} \bar{u} = -\frac{\ell^2}{r_+} e^{-u \frac{r_+}{\ell^2}}, \\ \bar{v} = +\frac{\ell^2}{r_+} e^{+v \frac{r_+}{\ell^2}}, \end{cases} \quad (17)$$

that results in the metric

$$ds^2 = \frac{(r + r_+)^2}{\ell^2} d\bar{u} d\bar{v} - r^2 d\phi^2, \quad (18)$$

which is free of any singularity, in particular, at the horizon. The inverse transition is given by

$$\begin{cases} t = \frac{\ell^2}{2r_+} \ln \left[ -\frac{\bar{v}}{\bar{u}} \right], \\ r_* = \frac{\ell^2}{2r_+} \ln \left[ -\bar{u} \bar{v} \cdot \frac{r_+^2}{\ell^4} \right]. \end{cases} \quad (19)$$

Metric (18) is useful for analyzing the causal relations, especially, in the radial motion, since isotropic lines, i.e. light cones ( $ds^2 = 0$ ), correspond to lines parallel to axes of  $\bar{v}$  and  $\bar{u}$  ( $d\bar{v} = 0$  or  $d\bar{u} = 0$ ).

Second, we introduce the coordinate map for the geodesics completely confined behind the horizon in terms of  $\rho$  and phase  $\varphi_E$

$$\begin{cases} \bar{u} = \varkappa i \rho e^{i\varphi_E}, \\ \bar{v} = -\frac{i}{\varkappa} \rho e^{-i\varphi_E}, \end{cases} \quad (20)$$

which are related with the primary time and radius by

$$\begin{cases} t = -i \frac{\ell^2}{r_+} \varphi_E + \Delta t_0, \quad \varphi_E \in [0, 2\pi], \\ r_* = \frac{\ell^2}{r_+} \ln \left[ -\frac{\rho r_+}{\ell^2} \right], \quad \rho \in [0, \ell^2/r_+], \end{cases} \quad (21)$$

at  $\Delta t_0 = -\ell^2/r_+ \ln \varkappa$ . Eq. (21) certainly establishes the value of period in the purely imaginary time, that determines the inverse temperature of black hole at  $J = 0$ ,

$$\beta_0 = 2\pi \frac{\ell^2}{r_+}. \quad (22)$$

The horizon ‘area’ (perimeter) is given by

$$\mathcal{A}_0 = 2\pi r_+ = 4\pi \ell \sqrt{2GM}, \quad (23)$$

where we have substituted  $r_+$  in terms of black hole mass at  $J = 0$ . Then, by differentiation we find

$$dM = \frac{T_0}{4G} d\mathcal{A}_0, \quad T_0 = \frac{1}{\beta_0}, \quad (24)$$

that indicates the introduction of entropy by

$$\mathcal{S}_0 = \frac{\mathcal{A}_0}{4G}, \quad (25)$$

in accordance with the Bekenstein–Hawking relation [5, 7, 8].

In order to calculate the partition function of particles moving by classical paths completely confined behind the

horizon [1], we, first, evaluate the increment of phase per cycle, i.e. by motion from  $r = 0$  to  $r = r_c$  and back,

$$\Delta_c \varphi_E = i \cdot 2 \int_0^{r_c} dr \frac{dt}{dr} \frac{r_+}{\ell^2} = 2 \int_0^{r_c} dr \frac{r_+}{r_+^2 - r^2} \sqrt{\frac{r_+^2 - r_c^2}{r_c^2 - r^2}}. \quad (26)$$

Introducing

$$\xi = r^2,$$

we can represent integral (26) by

$$\Delta_c \varphi_E = r_+ \oint \frac{d\xi}{2\sqrt{\xi}} \frac{1}{\xi_c - \xi} \sqrt{\frac{\xi_+ - \xi_c}{\xi_c - \xi}}, \quad (27)$$

where the contour of integration loops the cut in complex plane of  $\xi$  because of square roots in the integrand as shown in Fig. 1.




FIG. 1: The integration contour in complex plane of  $\xi$ .

However, we can consider the auxiliary contour including the pole at  $\xi_+$  in Fig. 2.




FIG. 2: The auxiliary contour in complex plane of  $\xi$ .

This contour can be deformed to the circle at infinity, where the integrand rather rapidly nullifies. Therefore, the integral over the auxiliary contour is equal to zero, which can be written as the sum of integral for the increment and integral around the pole. Hence, the increment of phase is given by the residue at  $\xi = \xi_+$ ,

$$\Delta_c \varphi_E = \pi. \quad (28)$$

Remarkably, it is independent of  $r_c$ .

In [1, 2, 4] we have introduced a winding number, determining a number of cycles per the full period of phase. By [4] we recognize that the increment value of (28) leads to the following conclusion: there is the only quantum level of particle moving behind the horizon of BTZ black

hole at  $J = 0$ , i.e. the level at  $r_c \rightarrow r_+$  or  $A \rightarrow -\infty$ , since at  $r_c < r_+$  we expect the winding number equal to  $3\pi/(2\Delta_c \varphi_E) = 3/2$ , which is not integer. The level is the ground state of particle in the thermal ensemble. The corresponding winding number is given by

$$n = \frac{2\pi}{\Delta_c \varphi_E} = 2. \quad (29)$$

Thus, each particle can occupy two orbits at the ground level.

The increment of interval is determined by

$$\Delta_c s_0 = 2 \int_0^{r_c} dr \frac{\ell}{\sqrt{r_c^2 - r^2}} = \pi \ell, \quad (30)$$

which is also independent of  $r_c$ . Therefore, the action of massive particle confined behind the horizon is equal to

$$S_0 = -mc n \Delta_c s_0 = -mc 2\pi \ell. \quad (31)$$

Following [1, 2], we deduce the thermodynamical function

$$G = -\beta \mathcal{F} = -2\pi \ell \sum mc, \quad (32)$$

where  $\mathcal{F}$  is the Helmholtz free energy. Further, we denote the sum of particle masses by

$$\sigma_0 = \sum mc. \quad (33)$$

Then, in the framework of thermodynamical relations, we fix the total energy of the system by the black hole mass

$$M = -\frac{\partial G}{\partial \beta} = 2\pi \ell \frac{\partial \sigma_0}{\partial \beta_0}. \quad (34)$$

However, at  $J = 0$  we can easily use

$$\frac{\partial \beta_0}{\partial M} = -\frac{\pi \ell}{2M} \frac{1}{\sqrt{2GM}},$$

so that

$$\frac{\partial \sigma_0}{\partial M} = -\frac{1}{4\sqrt{2GM}},$$

that gives

$$\sigma_0 = -\frac{1}{2} \sqrt{\frac{M}{2G}}, \quad (35)$$

where we have put the integration constant equal to zero, as usually accepted. Then, the entropy can be evaluated by standard formula of thermodynamics,

$$\mathcal{S}_0 = \beta_0 M + G = \pi \ell \sqrt{\frac{2M}{G}} = \frac{1}{4G} \mathcal{A}_0. \quad (36)$$

The only unusual feature of such the derivation is the negative sign of function  $\sigma_0$  equal to the sum of particle

masses. This fact is closely connected with the notion of vacuum for the black hole in the anti-de Sitter space (AdS). As was shown in [6, 9], the true vacuum is the AdS space corresponding to the mass parameter equal to  $M_{\text{AdS}} = -(8G)^{-1}$ , i.e. the negative value of mass. This vacuum is treated as a ‘bound’ state. By analogy, the thermodynamical ensemble has the negative free energy  $\mathcal{F}$  at  $J = 0$ ,

$$\mathcal{F}_0 = -M,$$

that could be treated as the indication of binding the particles in the field of BTZ black hole. If

$$\mathcal{F}_0 \geq M_{\text{AdS}} \Rightarrow M \leq \frac{1}{8G}, \quad r_+ \leq \ell, \quad (37)$$

hence, the horizon has a radius less than the curvature radius of AdS space.

The value of entropy is in agreement with calculations by A. Strominger in the framework of conformal theory [9]. Indeed, the entropy counts the number of states with the mass  $M$ , while the Virasoro algebra has the central charge

$$c = \frac{3\ell}{2G}, \quad (38)$$

and generators

$$L_0 = \bar{L}_0 = \frac{2M}{\ell}, \quad (39)$$

so that the Cardy formula [10] gives

$$\mathcal{S}_0 = 4\pi\sqrt{\frac{cL_0}{6}} = \pi\ell\sqrt{\frac{2M}{G}}. \quad (40)$$

The variation of  $G$  can be written in the form

$$dG = \frac{\pi\ell}{2\sqrt{2GM}} dM = \beta dM, \quad (41)$$

or

$$dG = -2\pi\ell d\sigma, \quad (42)$$

which are consistent with each other. The Hawking radiation takes place at  $dM < 0$ , i.e. at the decrease of black hole mass. Eq. (41) means that the probability distribution versus the energy emitted by the black hole  $E = -dM$ , is given by

$$w \sim e^{-\beta E}, \quad (43)$$

which is the Gibbs distribution, that reproduces the thermal spectrum of black body up to ‘grey body’ factors of re-scattering. The radiation is balanced by the appropriate change of particle masses at the ground level as dictated by (41) and (42),

$$d\sigma = -\frac{\beta}{2\pi\ell} dM.$$

The consideration of Hawking radiation as a quasi-classical tunnel effects was given in [11]. Further details could be also found in review on the thermodynamics of black hole horizons [12].

### III. TWO HORIZONS: $J \neq 0$

At  $J \neq 0$  we follow the consideration in second reference of [2]. So, we put the angle dependent contribution in the BTZ metric of (2) equal to zero,

$$\dot{\phi} = \frac{4JG}{\rho^2}, \quad (44)$$

which corresponds to a definite relation between conserved total energy  $E$  and orbital momentum  $\mu$ . Then, we get the interval

$$ds^2 = N^2 dt^2 - \frac{1}{N^2} d\rho^2 \quad (45)$$

in a strong analogy with the case of  $J = 0$ . The difference is due to the presence of two horizons and two return point for geodesics completely confined behind the horizons.

The interval of trajectory takes the form

$$ds^2 = \frac{d\rho^2}{1/A - N^2}, \quad (46)$$

where again  $A = m^2/E^2$ . For the confined geodesics  $A < 0$ , and the return points  $\rho_{1,2}$  are positive solutions of equation

$$\rho^4 - \left(8GM\ell^2 + \frac{\ell^2}{A}\right)\rho^2 + 16J^2G^2\ell^2 = 0. \quad (47)$$

With

$$\mathcal{B} = 1 + \frac{1}{8GMA}$$

roots are

$$\rho_{1,2}^2 = 4GM\ell^2 \left[ \mathcal{B} \pm \sqrt{\mathcal{B}^2 - \frac{J^2}{M^2\ell^2}} \right], \quad (48)$$

that indicates the existence of minimal value of negative  $A$ . The roots are arranged, so that

$$\rho_- \leq \rho_1 \leq \rho_2 \leq \rho_+,$$

of course.

The increment of interval is determined by

$$\Delta cs = 2 \int_{\rho_1}^{\rho_2} d\rho \frac{\ell\rho}{\sqrt{(\rho_2^2 - \rho^2)(\rho^2 - \rho_1^2)}} = \pi\ell. \quad (49)$$

It is fixed and independent of return points.

The ‘tortoise’ coordinate is given by

$$\begin{aligned} \rho_* &= \int d\rho \frac{\ell^2 \rho^2}{(\rho^2 - \rho_+^2)(\rho^2 - \rho_-^2)} = \\ &= \frac{\ell^2}{2(\rho_+^2 - \rho_-^2)} \left[ \rho_+ \ln \frac{\rho - \rho_+}{\rho + \rho_+} - \rho_- \ln \frac{\rho - \rho_-}{\rho + \rho_-} \right], \end{aligned} \quad (50)$$

so that, after the introduction of appropriate Kruskal coordinates and corresponding two maps with purely imaginary time [2], we find two inverse temperatures of external and inner horizons, correspondingly,

$$\beta_{\pm} = 2\pi \frac{\ell^2 \rho_{\pm}}{\rho_+^2 - \rho_-^2}. \quad (51)$$

The consistency of two maps takes place, if

$$\frac{\beta_+}{\beta_-} = \frac{\rho_+}{\rho_-} = k, \quad (52)$$

where  $k$  is an integer or half-integer number [3].

The increment of phase defined by the purely imaginary time,

$$\varphi_E = \frac{2\pi}{\beta_+} t_E, \quad (53)$$

is given by the integral

$$\Delta_c \varphi_E = \frac{2\pi}{\beta_+} 2 \int_{\rho_1}^{\rho_2} d\rho \frac{\sqrt{-1/A}}{N^2 \sqrt{1/A - N^2}}, \quad (54)$$

which can be evaluated by the same way of previous section, i.e. the contour integration after the transformation of variables  $\xi = \rho^2$ . Then, it is reduced to contribution of two residues by poles at  $\rho^2 = \rho_{\pm}^2$ , where  $N^2$  nullifies,

$$\Delta_c \varphi_E = \pi \frac{\rho_+ - \rho_-}{\rho_+}. \quad (55)$$

The winding number of ground level is equal to

$$n_+ = \frac{2\pi}{\Delta_c \varphi_E} = \frac{2k}{k-1}. \quad (56)$$

It should be integer, that restricts values of  $k$  by the following set:

$$k = \left\{ 1, \frac{3}{2}, 2, 3, \infty \right\}. \quad (57)$$

$k = 1$  represents the extremal black hole, while  $k = \infty$  reproduces the black hole with  $J = 0$ .

As we have already recognized in the previous section, there are no excitations of ground state for the particle confined behind the horizons of BTZ black hole.

Next, the action of single particle at the ground level is equal to

$$S = -mc n_+ \Delta_c s = -mc 2\pi \frac{\ell \rho_+}{\rho_+ - \rho_-}, \quad (58)$$

or

$$S = -mc \frac{\rho_+ + \rho_-}{\ell} \beta_+,$$

which gives the partition function

$$\ln Z = -\beta_+ \sigma \frac{\rho_+ + \rho_-}{\ell}, \quad (59)$$

where  $\sigma = \sum mc$  is the sum of particle masses ( $c = 1$ ). Following the method of [2], the total energy of system is equal to the mass of black hole,

$$M = -\frac{\partial \ln Z}{\partial \beta_+},$$

if

$$\sigma = \frac{\ell}{\rho_+ + \rho_-} \left( M - \frac{1}{4G\beta_+} \mathcal{A}_+ \right), \quad (60)$$

which reproduces the result of previous section for the sum of masses at  $\rho_+ = r_+$  and  $\rho_- = 0$ .

Further, in the standard thermodynamical way, the entropy

$$\mathcal{S} = \beta_+ M + \ln Z$$

is equal to

$$\mathcal{S} = \frac{1}{4G} \mathcal{A}_+, \quad (61)$$

in agreement with the Bekenstein–Hawking formula.

The Helmholtz energy

$$\mathcal{F} = M - \frac{\mathcal{A}_+}{4G\beta_+} = M \left( 1 - 2\sqrt{1 - \frac{J^2}{M^2 \ell^2}} \right)$$

is positive at

$$J^2 \geq \frac{3}{4} M^2 \ell^2 \Rightarrow \mathcal{F} \geq 0. \quad (62)$$

#### IV. QUANTUM SPECTRUM

The relation between the BTZ black hole mass  $M$  and its orbital momentum  $J$  can be easily obtained from the condition of map consistency (52),

$$\frac{\rho_+^2}{\rho_-^2} = k^2 \Leftrightarrow k^2 = \frac{1 + \sqrt{1 - \frac{J^2}{M^2 \ell^2}}}{1 - \sqrt{1 - \frac{J^2}{M^2 \ell^2}}}, \quad (63)$$

that gives<sup>1</sup>

$$J = \frac{2k}{k^2 + 1} M \ell, \quad (64)$$

which indicates the linear dependence of mass  $M$  versus the orbital momentum  $J$ . Then, the set of  $k$  values determines 5 quasi-classical trajectories in the plane of  $\{J, M\}$ .

The set of winding numbers

$$n_+ = \{\infty, 6, 4, 3, 2\}, \quad (65)$$

---

<sup>1</sup> Here we put  $J \geq 0$ , for definiteness.

can be treated as numbers of massive spin-states:  $n_+ = 2s + 1$  (at  $J \neq 0$  and in non-extremal case).

Next, the ‘area’ is also quantized as

$$\mathcal{A}_+ = 2\pi\rho_+ = 4\pi\sqrt{G\ell Jk}. \quad (66)$$

It is spectacular that the ‘area’ is not linear in integer or half-integer  $J$ .

## V. CONCLUSION

We have demonstrated that the approach of quasi-classical quantization for the particles completely con-

fined behind the horizon [1, 2, 3, 4] can be successfully applied to the BTZ black hole. The features of BTZ black hole are the followings: the absence of excitations over the ground level, and the linear dependence of mass versus the orbital momentum.

We could expect that the ground level of particles in the BTZ black hole would be quite a coherent state, whose characteristics at the external horizon is complete in a sense of holographic principle [13].

This work is partially supported by the grant of the president of Russian Federation for scientific schools NSC-1303.2003.2, and the Russian Foundation for Basic Research, grant 04-02-17530.

- 
- [1] V. V. Kiselev, arXiv:gr-qc/0412090.
  - [2] V. V. Kiselev, arXiv:gr-qc/0412097, gr-qc/0412110,
  - [3] V. V. Kiselev, arXiv:gr-qc/0412117.
  - [4] V. V. Kiselev, arXiv:gr-qc/0506006.
  - [5] S. W. Hawking, Commun. Math. Phys. **43**, 199 (1975);  
J. B. Hartle and S. W. Hawking, Phys. Rev. D **13**, 2188 (1976).
  - [6] M. Banados, C. Teitelboim and J. Zanelli, Phys. Rev. Lett. **69**, 1849 (1992) [arXiv:hep-th/9204099];  
M. Banados, M. Henneaux, C. Teitelboim and J. Zanelli, Phys. Rev. D **48**, 1506 (1993) [arXiv:gr-qc/9302012].
  - [7] J. D. Bekenstein, Lett. Nuovo Cim. **4**, 737 (1972);  
J. D. Bekenstein, Phys. Rev. D **7**, 2333 (1973);  
J. D. Bekenstein, Phys. Rev. D **9**, 3292 (1974).
  - [8] S. W. Hawking, Phys. Rev. D **13**, 191 (1976);  
G. W. Gibbons and S. W. Hawking, Phys. Rev. D **15**, 2738 (1977);  
S. W. Hawking and G. T. Horowitz, Class. Quant. Grav. **13**, 1487 (1996) [arXiv:gr-qc/9501014].
  - [9] A. Strominger, JHEP **9802**, 009 (1998) [arXiv:hep-th/9712251].
  - [10] J. L. Cardy, Nucl. Phys. B **270**, 186 (1986);  
H. W. J. Bloete, J. L. Cardy and M. P. Nightingale, Phys. Rev. Lett. **56**, 742 (1986).
  - [11] E. Keski-Vakkuri and P. Kraus, Phys. Rev. D **54**, 7407 (1996) [arXiv:hep-th/9604151];  
M. K. Parikh and F. Wilczek, Phys. Rev. Lett. **85**, 5042 (2000) [arXiv:hep-th/9907001];  
M. Angheben, M. Nadalini, L. Vanzo and S. Zerbini, arXiv:hep-th/0503081;  
E. C. Vagenas, Phys. Lett. B **559**, 65 (2003) [arXiv:hep-th/0209185];  
E. C. Vagenas, Phys. Lett. B **533**, 302 (2002) [arXiv:hep-th/0109108];  
E. C. Vagenas, Mod. Phys. Lett. A **17**, 609 (2002) [arXiv:hep-th/0108147];  
M. Arzano, A. J. M. Medved and E. C. Vagenas, arXiv:hep-th/0505266;  
K. Srinivasan and T. Padmanabhan, Phys. Rev. D **60**, 024007 (1999) [arXiv:gr-qc/9812028];  
T. Padmanabhan, Mod. Phys. Lett. A **19**, 2637 (2004) [arXiv:gr-qc/0405072].
  - [12] T. Padmanabhan, Phys. Rept. **406**, 49 (2005) [arXiv:gr-qc/0311036].
  - [13] G. 't Hooft, arXiv:gr-qc/9310026;  
L. Susskind, J. Math. Phys. **36**, 6377 (1995) [arXiv:hep-th/9409089];  
R. Bousso, Rev. Mod. Phys. **74**, 825 (2002) [arXiv:hep-th/0203101].